



# BIOTECHNOLOGY *for* A CIRCULAR BIOECONOMY

28 - 29 March 2023 | AFOB-EFB Virtual Conference

## Programme and abstract book

Platinum  
Sponsor



**ELSEVIER**

**Biotechnology for a circular  
bioeconomy:  
carbon capture, waste recycling  
and mitigation of global warming**

## TABLE OF CONTENTS

PROGRAMME.....	5
LIST OF FLASH POSTER PRESENTATIONS.....	9
ABSTRACTS .....	11
<b>Oral presentations .....</b>	<b>12</b>
Session 1: Carbon capture biotechnology: from CO <sub>2</sub> to protein.....	13
Session 2: From waste and wastewater to valuable products.....	20
Session 3: Biotechnology to mitigate climate change and global warming.....	26
Session 4: Towards a circular bioeconomy: progress and bottlenecks .....	33
<b>Poster presentations.....</b>	<b>39</b>
INDEX OF AUTHORS.....	104



europaean  
federation of  
biotechnology



ELSEVIER

## **A biotechnological process for the production of pyocyanin and 1-hydroxyphenazine using waste streams from the potato chips industry**

Lena Pantelic, Jelena Lazic, Jasmina Nikodinovic-Runic, Sandra Vojnovic

Institute of Molecular Genetics and Genetic Engineering

Potato industry is one of the major food industries that generates considerable amounts of potato peels and wastewater (1). These wastes can then be utilized as components of microbial media in biotechnological production of pyocyanin (PYO) and 1-hydroxyphenazine (1-HP) using *Pseudomonas aeruginosa* (2). PYO and 1-HP possess important biological activities, thus could be applied in the field of medicine, and can be used as biocontrolling agents (2). However, their application is hindered due to high costs associated with their large scale production.

In this work, we established a fermentation process which utilizes either potato peels or potato wastewater as the sole nutrient source to obtain PYO and 1-HP. *P. aeruginosa* BK25H strain was selected from our in-house collection. This approach afforded 10 mg/l PYO and 9 mg/l 1-HP using potato wastewater and 15 mg/l PYO and 11 g/l 1-HP using potato peels after 24 h incubation. This work is the step towards zero-pollution and conversion of waste to valuable microbial products.

### References

(1) Radeef A.Y. and Ismail Z.Z. *Int. J. Green Energy* 2021, 18, 910.

(2) Pantelic L., Skaro Bogojevic S., Vojnovic S., Lazic J., Ilic-Tomic T., Milivojevic D., Nikodinovic-Runic J. *Enzyme Microb. Technol.* 2023 (unpublished)

Acknowledgments - This research was funded by the Science Fund of the Republic of Serbia, IDEAS, 7730810, BioECOLogics. The potato peels and potato wastewater were kindly supplied by Polo Ltd. (Cacak, Serbia).

**efb** | european  
federation of  
**biotechnology**

